

## **Consent for Botulinum Toxins**

(Botox, Dysport, Nuceiva)

Botulin um toxin (B-tox) therapy for wrinkles is an injection treatment designed to reduce facial expression lines. B-tox is approved for the treatment of disorders of the muscles of the eye, certain types of headaches and other medical conditions. As well, it is approved for cosmetic treatment of frown lines between and just above the eyebrows. B-tox therapy for all such wrinkles is a commonly performed cosmetic procedure throughout the world.

When small amounts of purified B-tox are injected into a muscle, contractions are greatly diminished or shut down completely. The result is fewer wrinkles in the overlying skin. Medically approved areas of the face include the frown lines (corrugators and procerus muscles), horizontal forehead lines (frontalis) and the crow's feet (periorbital muscles).

As well, B-tox is frequently used "off-label" in other areas of the face including the chin and lip regions, bands about the neck, and bunny lines above the nose. A small percentage of clients will notice results the very next day, however the majority will appreciate the change within 3-7 business days. A small percentage may not notice optimal results before 2 weeks. The results typically last 3-4 months but be shorter or longer.

For reasons not understood, a small percentage of people notice the treatment effects vary from one treatment to the next. Neutralizing antibodies (proteins) produced by your body may reduce the effectiveness of this treatment. The potential for such formation of such antibodies can be minimized by using the lowest effective dose and spreading out the treatments as best one can.

1. I understand that I will not be able to move the treated muscles while the injection is

## Results, Risks and Aftercare

<ul> <li>2. I understand that during the 4-hour post-injection period, forward (sitting upright such as while driving is fine) or exarea of the injection site</li> <li>3. I understand the following possible temporary side effects</li> <li>Bruising is very common, and the risk may increase anti-inflammatories such as Aspirin, Ibuprofen and</li> </ul>	it is bost not to lie down bond
<ul> <li>area of the injection site</li> <li>I understand the following possible temporary side effects</li> <li>Bruising is very common, and the risk may increase</li> </ul>	•
Bruising is very common, and the risk may increase	ercise, or manipulate/rub the
, , ,	s:
<ul> <li>such as Vitamin E, Omega III's (fish oils), Ginkgo Bil Ginger or Cayenne, as well as Alcoholic Beverages. temporary, will resolve in 5-10 days, and will not a treatment.</li> <li>I understand that a headache or pain at the injecti is a temporary effect lasting less than 24 hours. Ac be taken for this after the treatment.</li> </ul>	d Naproxen, or supplements iloba, St. John's Wort, Garlic, I understand that this is affect the outcome of the ion site may occur but that this

•				s after treatment. Usually p" if unresolved after such time		
•	<ul> <li>Numbness or dullness, or change in sensation to the treated areas moticed for a few days after the injection. Treatment for this is not need to be a few days.</li> </ul>					
•	an eyelid (ptosis) rarely up to severa Double vision (dip temporary.  I understand that satisfactorily or fo	treatment of frow in 1-2% of frown li al months, but can olopia) may occur vor a very small nur as long as expect	n lines may rene injections.  be treated with crow's featings.  Imber of pations.	esult in a temporary drooping of This can last 2-3 weeks and with eye dropset injections. This is also ents, Botox does not work as		
immur	ot aware of having nosuppressant med ot aware that I am	lications				
I have read th		stand. My questio	ns have been	answered to my satisfaction by		
Name (please	print)		<u>-</u>			
Patient Signa	iture	MD Signature		 Date		